



Bouncy Concealer

Bouncy Concealer with Creaspheres® EL WL 7 which gives unique cushioning sensation and flexibility to the formulation. Eospolys® are UV filters where the Titanium Dioxide is encapsulated in Silica bead or polymer structure. BNPoly® UV Crystals boost the UV protection of the formulation and create ceramic effect upon application. Creasil® 11 IH is a film former that provides long lasting water resistance, but transparent and smooth film. Novatext® provides elegant, silicone-like texture and reduces any tackiness in the formulation. Alphaflow® and Dedraflow® are photostable and hypoallergenic emollients. Dedraflow® 5 is volatile and can be used as replacement to Cyclomethicone. Hectone® is an ideal thickener, which prevents any risk of sedimentation. Creasperse® Colours are stable colour dispersions. Fiflow® is an anti-age active giving instant visual results. Fiflow® reduces the appearance of fine lines as it works as a muscle relaxant and wrinkle filler.

Ingredients	INCI Name	Qty%	Supplier
Phase A			
Eospoly® TR22 SIL 30	Silica (and) Titanium Dioxide (and) Alumina	9,78	1)
Creaspheres® EL WL7	Vinyl Dimethicone/Methicone Silsesquioxane Crosspolymer (and) Polymethylsilsesquioxane	7,65	1)
Talcpoly® WL4	Talc	7,65	1)
BNPoly® UV Crystal TR 14	Boron Nitride (and) Titanium Dioxide (and) Dimethicone (and) Isododecane (and) Ethylene/VA Copolymer	14,25	1)
Phase B			
Creasil® 11IH	Isohexadecane (and) Dimethicone (and) Polyethacrylate	6,40	1)
Creasil® IHCG	Isohexadecane	2,35	1)
Novatext® Velvet	Hydrogenated Polyisobutene (and) Polymethylsilsesquioxane (and) Vinyl Dimethicone/Methicone Silsesquioxane Crosspolymer (and) Ethylene/Propylene Copolymer	5,86	1)
Alphaflow® 20	Hydrogenated Polydecene	2,75	1)
Alphaflow® 50	Hydrogenated Polydecene	2,31	1)
Dedraflow® 325	Hydrogenated Polyisobutene (and) Dimethicone	6,34	1)
Dedraflow® 5	Hydrogenated Polyisobutene	6,67	1)
Hectone® DF	Hydrogenated Polyisobutene (and) Distearidimonium Hectorite (and) Propylene Carbonate	4,05	1)
Creasperse® Honey Beige	Hydrogenated Polydecene (and) CI77492 (and) CI77891 (and) Hydroxystearic Acid (and) CI77491 (and) CI77499	9,17	1)
Creasperse® White R	Titanium Dioxide (and) Hydrogenated Polydecene (and) Hydroxystearic Acid	12,00	1)
Phase C			
Fiflow® SC	Perfluorohexane (and) Perfluorodecalin (and) Perfluoroperhydrophenanthrene (and) Perfluorodimethylcyclohexane	2,77	1)

Procedure:

1. Keep phase A under agitation until homogeneous. No heating necessary.
2. Keep phase B under agitation until homogeneous.
3. Add phase B into phase A and keep under agitation until homogeneous.
4. Add phase C into the mixture (A+B) and keep under agitation until homogeneous.

Suppliers:

- 1) CIT SARL